## For Research Use Only

## Histone H1.5 Polyclonal antibody Catalog Number: 18093-1-AP





| Basic Information      | Catalog Number:<br>18093-1-AP   | GenBank Accession Number<br>BC101581 | : Purification Method:<br>Antigen affinity purification   |
|------------------------|---|--------------------------------------|---|
|                        | Size:<br>260 ug/ml  | GenelD (NCBI):<br>3009               | Recommended Dilutions:<br>WB 1:5000-1:30000   |
|                        | Source:<br>Rabbit   | UNIPROT ID:<br>P16401                | IHC 1:50-1:500<br>IF/ICC 1:50-1:500   |
|                        | Isotype:<br>IgG   | Full Name:<br>histone cluster 1, H1b |   |
|                        | Immunogen Catalog Number:<br>AG12818  | Calculated MW:<br>226 aa, 23 kDa     |   |
|                        |   | Observed MW:<br>32 kDa               |   |
| Applications           | WB, IHC, IF/ICC, ELISA WB : He   Species Specificity: IHC : m   human, mouse colon ti   Note-IHC: suggested antigen retrieval with  |                                      | ive Controls:   |
|                        |   |                                      | HeLa cells, HepG2 cells, LnCaP cells<br>mouse brain tissue, human colon tissue, mouse<br>n tissue |
|                        |   |                                      | C : SH-SY5Y cells,  |
| Background Information | Histones are basic nuclear proteins responsible for nucleosome structure of the chromosomal fiber in eukaryotes.<br>Histone H1.5, also known as Histone Cluster 1 H1 Family Member B (HIST1H1B), is a member of the Histone H1/H5<br>protein family. Histone H1.5 gene has been found to be frequently mutated in colorectal cancer and follicular<br>lymphomas. HIST1H1B overexpression contributes to tumor aggressiveness. Histone H1.5 expression is upregulated<br>in breast cancer and predicts poor prognosis(PMID: 34746019). |                                      |   |
| Storage                | Storage:<br>Store at -20°C. Stable for one year<br>Storage Buffer:<br>PBS with 0.02% sodium azide and<br>Aliquoting is unnecessary for -20°   | 50% glycerol pH 7.3.                 |   |

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 18093-1-AP (Histone H1.5 antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 18093-1-AP (Histone H1.5 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 18093-1-AP (Histone H1.5 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using Histone H1.5 antibody (18093-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human normal colon slide using 18093-1-AP (Histone H1.5 antibody) at dilution of 1:6000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).