For Research Use Only

NOP56 Polyclonal antibody

Catalog Number: 18181-1-AP 2 Publications



Basic Information

Catalog Number: GenBank Accession Number: 18181-1-AP BC104791 GeneID (NCBI): Size: 500 ug/ml 10528 **UNIPROT ID:** Source: Rabbit 000567

Full Name: Isotype:

NOP56 ribonucleoprotein homolog

(veast) Immunogen Catalog Number:

AG12882 Calculated MW:

594 aa, 66 kDa Observed MW: 66 kDa

Applications

Tested Applications: WB, IHC, ELISA **Cited Applications:**

Species Specificity: human, mouse, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000 IHC 1:50-1:500

Positive Controls:

WB: NIH/3T3 cells, A549 cells, HeLa cells

IHC: human stomach tissue,

Background Information

Box C/D RNA-protein complexes (RNPs) guide the 2'-O-methylation of nucleotides in eukaryotic ribosomal RNAs. The box C/D and C'/D' RNP subcomplexes are each assembled with three sRNP core proteins. The Nop56 core protein mediates crucial protein-protein interactions required for both sRNP assembly and the methyltransferase reaction by bridging the L7Ae and fibrillarin core proteins [PMID:22496443]. NOP56 is a component of the box C/D small nucleolar ribonucleoprotein complexes that direct 2-prime-O-methylation of pre-ribosomal RNA (rRNA) during its maturation. It is predicted to function in an early to middle step in pre-rRNA processing [PMID:12777385].

Notable Publications

Author	Pubmed ID	Journal	Application
Marlene Cervantes	32450515	iScience	WB
Miao Bai	34512161	Int J Biol Sci	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 18181-1-AP (NOP56 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 18181-1-AP (NOP56 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 18181-1-AP (NOP56 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).