

For Research Use Only

SHBG Polyclonal antibody

Catalog Number: 18200-1-AP

3 Publications



Basic Information

Catalog Number:

18200-1-AP

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG12683

GenBank Accession Number:

BC101785

GeneID (NCBI):

6462

UNIPROT ID:

P04278

Full Name:

sex hormone-binding globulin

Calculated MW:

402 aa, 44 kDa

Observed MW:

42-44 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:500-1:2000

IHC: 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse

Cited Species:

goat

Positive Controls:

WB : human plasma,

IHC : mouse testis tissue, human liver tissue

Note-IHC: suggested antigen retrieval with *TE buffer pH 9.0*; (*) Alternatively, antigen retrieval may be performed with *citrate buffer pH 6.0*

Background Information

SHBG (sex hormone-binding globulin), also known as androgen-binding protein (ABP), is the specific transport protein for sex steroid hormones in humans. SHBG is synthesized mainly by the liver and circulates in the blood, in form of homodimer with a single steroid-binding site per dimer. In addition to the plasma SHBG, a testis isoform expressed by Sertoli cells also exists due to the gene alternative splicing. Various bands (44-56 kDa) recognized by this antibody may represent variably glycosylated or different isoforms of SHBG.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|------------|-----------|---------------------|-------------|
| Huaming Xi | 34774699 | Mol Cell Endocrinol | WB |
| Huaming Xi | 36332369 | Theriogenology | WB |
| Huaming Xi | 34467369 | Biol Reprod | WB |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human plasma were subjected to SDS PAGE followed by western blot with 18200-1-AP (SHBG antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 18200-1-AP (SHBG antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 18200-1-AP (SHBG antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).