### For Research Use Only

# PHLDA1 Polyclonal antibody

Catalog Number: 18263-1-AP

Featured Product

9 Publications



**Basic Information** 

Catalog Number:

18263-1-AP

BC018929

Concentration:

600 ug/ml

22822

Source:

UNIPROT ID:

Rabbit

Q8WV24

Isotype:

GenBank Accession Number:

BC018929

GeneID (NCBI):

22822

UNIPROT ID:

Q8WV24

Full Name:

pleckstrin homology-like domain,

Immunogen Catalog Number: family A, member 1
AG13125 Calculated MW:
45 kDa

Observed MW: 40-45 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IF Species Specificity: human, mouse

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: A375 cells, mouse brain tissue, BXPC-3 cells, HeLa

**Purification Method:** 

WB 1:1000-1:6000 IHC 1:400-1:1600

IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

cells, MDA-MB-231 cells

IHC: human skin cancer tissue,

IF/ICC: A375 cells,

## Background Information

PHLDA1, also known as PHRIP and TDAG51, is a multifunctional protein involved in various biological processes. It can induce apoptosis in various cell types, including T cells, hippocampal cells, endothelial cells, melanoma cells, and mouse embryonic fibroblasts(PMID: 30207029). PHLDA1 plays a role in inhibiting growth factor signaling and has been implicated in tumor suppression through its ability to repress Akt activity by binding to phosphatidylinositol (PIP) lipids(PMID: 36142223).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
F Yang	33938317	Hum Exp Toxicol	WB
Liang Liu	31189920	Cell Death Dis	WB,IHC,IF
Yuxuan Guo	31981628	Life Sci	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 18263-1-AP (PHLDA 1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded skin cancer slide using 18263-1-AP (PHLDA1 antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed A375 cells using PHLDA1 antibody (18263-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).