For Research Use Only

Pinin Polyclonal antibody

Catalog Number: 18266-1-AP

Featured Product

7 Publications

GenBank Accession Number:



Purification Method:

WB 1:500-1:2000

IF/ICC 1:300-1:1200

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: 18266-1-AP

Size: 500 ug/ml BC062602 GeneID (NCBI): 5411

Source: UNIPROT ID:
Rabbit Q9H307
Isotype: Full Name:

gG pinin, desmosome associated protein

Immunogen Catalog Number: Calculated MW:

AG6643 82 kDa

Observed MW: 82 kDa, 140 kDa

Applications

Tested Applications: WB, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, RIP Species Specificity: human

Cited Species: human, mouse Positive Controls:

WB: HeLa cells, HepG2 cells

IP : HeLa cells,
IF/ICC : HeLa cells,

Background Information

PNN, also named as DRS or 140 kDa nuclear and cell adhesion-related phosphoprotein, is a 717 amino acid protein, which belongs to the pinin family. PNN localizes in the plasma membrane and is expressed in placenta, lung, liver, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon, heart, epidermis, esophagus, brain and smooth and skeletal muscle. PNN is expressed strongly in melanoma metastasis lesions and advanced primary tumors. PNN as a transcriptional activator binds to the E-box 1 core sequence of the E-cadherin promoter gene. PNN is involved in the establishment and maintenance of epithelia cell-cell adhesion. PNN is a potential tumor suppressor for renal cell carcinoma. The calculated molecular weight of PNN is 82 kDa (native form), but the post-modified protein is 140 kDa (phospho form of isoform 1).

Notable Publications

| Author | Pubmed ID | Journal | Application |
|--------------|-----------|------------|-------------|
| Xuejun Yang | 27175589 | Oncotarget | WB |
| Ting Tang | 32364663 | Mol Oncol | WB,IP,IF |
| Liyuan Zhang | 27384878 | Oncotarget | WB |

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18266-1-AP (Pinin antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Pinin antibody (18266-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Pinin antibody (18266-1-AP) at dilution of 1:600 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).



IP result of anti-Pinin (IP:18266-1-AP, 4ug; Detection:18266-1-AP 1:300) with HeLa cells lysate 520ug.