

For Research Use Only

BMF Polyclonal antibody

Catalog Number: 18298-1-AP

Featured Product

4 Publications



Basic Information

Catalog Number:

18298-1-AP

Size:

300 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC069505

GeneID (NCBI):

90427

UNIPROT ID:

Q96LC9

Full Name:

Bcl2 modifying factor

Calculated MW:

21 kDa

Observed MW:

24 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:800

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

human, mouse

Positive Controls:

WB : human placenta tissue, HEK-293 cells

Background Information

BMF belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities such as central regulators of apoptotic cell death. BMF contains a single BCL2 homology domain 3 (BH3), and has been shown to bind BCL2 proteins and function as an apoptotic activator. This protein is found to be sequestered to myosin V motors by its association with dynein light chain 2, which may be important for sensing intracellular damage and triggering apoptosis. Alternatively spliced transcript variants encoding different isoforms have been identified. BMF is widely expressed in tissues such as pancreas, liver and kidney, and in hematopoietic tissues.

Notable Publications

Author	Pubmed ID	Journal	Application
Yu Guohua G	23133614	PLoS One	WB
Kaiqin Ye	32620849	Nat Commun	WB
Feng Xu	35280673	Int J Biol Sci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human placenta tissue were subjected to SDS PAGE followed by western blot with 18298-1-AP (BMF antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.