For Research Use Only

Cytokeratin 10 Polyclonal antibody

Source: Rabbit

Isotype:

Catalog Number: 18343-1-AP 19 Publications



Basic Information

Catalog Number: 18343-1-AP

BC034697 GeneID (NCBI): 3858 UNIPROT ID:

GenBank Accession Number:

IgG P13645
Immunogen Catalog Number: Full Name:
AG13136 keratin 10

Calculated MW: 584 aa, 59 kDa Observed MW: 50-59 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: WB: 1:5000-1:20000 IHC: 1:500-1:2000 IF-P: 1:50-1:500

FC (Intra): 0.40 ug per 10^6 cells in a

100 µl suspension

IF/ICC: 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse skin tissue, A431 cells, rat skin tissue IHC: human cervical cancer tissue, human breast cancer tissue, human lung cancer tissue, human skin cancer tissue. rat skin tissue

IF-P: mouse skin tissue,
IF/ICC: A431 cells,
FC (Intra): A431 cells,

Background Information

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. As a type I keratin, keratin 10 is a suprabasal marker of differentiation in stratified squamous epithelia.

Notable Publications

Author	Pubmed ID	Journal	Application
Lisa Lemoine	34579573	mBio	IF
Yingzheng Zhao	36240893	Int J Biol Macromol	IF
Shidi Wu	31790685	Life Sci	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18343-1-AP (Cytokeratin 10 antibody) at dilution of 1:200000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 18343-1-AP (Cytokeratin 10 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 18343-1-AP (Cytokeratin 10 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse skin tissue using Cytokeratin 10 antibody (18343-1-AP) at dilution of 1:200 and Coralite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using Cytokeratin 10 antibody (18343-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1X10^6 A431 cells were intracellularly stained with 0.4 ug Anti-Human Cytokeratin 10 (18343-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:1000 (red), or 0.4 ug rabbit 1gG isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.