## For Research Use Only

# KLF13 Polyclonal antibody

Catalog Number: 18352-1-AP

Featured Product

14 Publications



**Basic Information** 

Catalog Number:

18352-1-AP

Size:

350 µg/ml

Source:

Rabbit

Q9Y2Y9

Isotype:

GenBank Accession Number:

BC012741

GeneID (NCBI):

51621

UNIPROT ID:

Q9Y2Y9

Full Name:

IgG Kruppel-like factor 13
Immunogen Catalog Number: Calculated MW:

AG13200 31 kDa

Observed MW: 37-45 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000 IHC 1:20-1:200

**Applications** 

Tested Applications: WB, IHC, ELISA Cited Applications: WB, IHC, IF Species Specificity: human, rat, mouse Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HT-29 cells, rat liver tissue IHC: human brain tissue,

## **Background Information**

Kruppel-like factor 13(KLF13) is a member of transcription actors, which contains 3 zinc finger DNA-binding domains. Theses transcription factors bind to CG-rich sequences and related GT and CACCC boxes. KLF13 involved in activation of transcription of variety of genes in the cells of erythroid lineage, such as GATA1, glycophorin B. It also binds to the A and A/B RANTES promoter binding sites, the same as NFKB, and then transactivates the RANTES gene. The calculated molecular weight of KLF13 is 31 kDa, but the modified KLF13 protein is about 37-45 kDa.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Joshua E Burda	35614216	Nature	IHC
Chao-Chun Chen	36381108	J Clin Transl Hepatol	WB,IHC
Weilong Yao	32523679	Cell Biosci	WB

### Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

# Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 18352-1-AP (KLF13 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human brain using 18352-1-AP (KLF13 antibody) at dilution of 1:50 (under 10x lens).