

For Research Use Only

NOL7 Polyclonal antibody

Catalog Number: 18424-1-AP

Featured Product



Basic Information

Catalog Number:

18424-1-AP

Concentration:

300 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG13272

GenBank Accession Number:

BC023517

GeneID (NCBI):

51406

UNIPROT ID:

Q9UMY1

Full Name:

nucleolar protein 7, 27kDa

Calculated MW:

257 aa, 29 kDa

Observed MW:

38 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB : HEK-293 cells, HeLa cells

Background Information

NOL7 (nucleolar protein 7) is localized in both the nucleolus and nucleoplasm, with its nucleolar localization being RNA-dependent. Its function in the nucleolus is likely related to maintaining nucleolar structure and cell growth, while its role in the nucleoplasm may be associated with anti-angiogenic effects. Additionally, NOL7 is involved in the accumulation of early pre-rRNA and the processing of pre-18S rRNA, which are crucial for maintaining nucleolar function and cell growth. The expression of NOL7 is regulated by various transcription factors, including c-myc and RXR α . In cervical cancer, NOL7 transcription is also positively regulated by the RB tumor suppressor gene. The NOL7 gene encodes three distinct splice variants, each with different intracellular localizations that may affect their functions.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



WB result of NOL7 antibody (18424-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NOL7 transfected HEK-293 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 18424-1-AP (NOL7 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.