

For Research Use Only

PDLIM4 Polyclonal antibody

Catalog Number: 18543-1-AP



Basic Information

Catalog Number: 18543-1-AP	GenBank Accession Number: BC016765	Purification Method: Antigen affinity purification
Size: 350 ug/ml	GeneID (NCBI): 8572	Recommended Dilutions: WB 1:1000-1:6000
Source: Rabbit	UNIPROT ID: P50479	IHC 1:300-1:1200
Isotype: IgG	Full Name: PDZ and LIM domain 4	IF/ICC 1:200-1:800
Immunogen Catalog Number: AG13424	Calculated MW: 35 kDa	
	Observed MW: 35 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls:
Species Specificity: human	WB : A431 cells, K-562 cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : human skin cancer tissue,
	IF/ICC : U2OS cells,

Background Information

PDLIM4, also known as RIL, is a member of the ALP/Enigma family. PDLIM4 contains one PDZ(DHR) domain and one LIM zinc-binding domain. The LIM zinc-binding domain interacts with the second and fourth PDZ domains of Fap-1, possibly resulting in the dephosphorylation of PDLIM4 (PMID: 21636573). PDLIM4 is involved in osteoblast development and function. The variation of the PDLIM4 locus may be an important determinant of osteoporosis (PMID: 12908099).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 18543-1-AP (PDLIM4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded skin cancer slide using 18543-1-AP (PDLIM4 antibody) at dilution of 1:600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using PDLIM4 antibody (18543-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).