For Research Use Only

LSS Polyclonal antibody

Catalog Number: 18693-1-AP 3 Publications



Basic Information

Catalog Number: GenBank Accession Number: 18693-1-AP BC035638
Size: GeneID (NCBI): 4047
Source: UNIPROT ID: Rabbit P48449

Rabbit P48449

Isotype: Full Name:
IgG lanosterol synthase (2,3-

Immunogen Catalog Number: oxidosqualene-lanosterol cyclase)
AG6955 Calculated MW:

6955 Calculated MW: 732 aa, 83 kDa

Observed MW: 70-75 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

protein lysate IHC 1:50-1:500 IF/ICC 1:20-1:200

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WR IF

Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HeLa cells, IP: HeLa cells,

IHC: human liver cancer tissue,

IF/ICC: HepG2 cells,

Background Information

LSS, also named as OSC, is lanosterol synthase which catalyzes the cyclization of (S)-2,3 oxidosqualene to lanosterol, a reaction that forms the sterol nucleus. LSS gene has three isoforms with MW 74 kDa and 82-85 kDa. This antibody can recognize all these three isoforms.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiao-Yi Lu	33177714	Nature	WB
Minglei Zhao	34926465	Front Cell Dev Biol	WB,IF
Mei-Yan Ma	39489939	Cell Rep	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 18693-1-AP (LSS antibody) at dilution of 1:2000 incubated at room temperature for 1.5 burs.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 18693-1-AP (LSS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-LSS (IP:18693-1-AP, 4ug; Detection:18693-1-AP 1:500) with HeLa cells lysate 1880 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 18693-1-AP (LSS antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit lgG(H+L).