

For Research Use Only

ZBTB7A Polyclonal antibody

Catalog Number: 18700-1-AP

1 Publications



Basic Information

Catalog Number:

18700-1-AP

Concentration:

600 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC113511

GeneID (NCBI):

51341

UNIPROT ID:

O95365

Full Name:

zinc finger and BTB domain containing 7A

Calculated MW:

61 kDa

Observed MW:

61-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

human, mouse

Positive Controls:

WB : A549 cells, HEK-293T cells, NCI-H1299 cells, U2OS cells

IF/ICC : MCF-7 cells,

Background Information

ZBTB7A, also named as LRF, FBI1, FBI-1, pokemon, ZBTB7, ZNF857A and DKFZp5470146, a member of the POK (POZ and Krüppel) family of transcription factors, plays a role in differentiation, oncogenesis, and adipogenesis (PMID: 35798238). ZBTB7A is involved in the instruction of early lymphoid progenitors to develop into B lineage by repressing T-cell instructive Notch signals. ZBTB7A is widely expressed in normal thymus and it is aberrantly overexpressed in human cancers (PMID: 28942243). ZBTB7A has 10 potential sumoylation sites located at lysine 61, 354, 371, 379, 383, 396, 486, 487, 536 and 539 (PMID: 20471975, 19471103). ZBTB7A has a predicted molecular weight of 61 kDa and the 75-80-kDa band could be the sumoylated form.

Notable Publications

Author	Pubmed ID	Journal	Application
Jing-Ping Zhou	31998789	Biomed Res Int	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 18700-1-AP (ZBTB7A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using ZBTB7A antibody (18700-1-AP) at dilution of 1:400 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4), CL488-phalloidin (green).