## For Research Use Only

## HOXD13 Polyclonal antibody Catalog Number: 18736-1-AP 3 Publications




| Basic Information      | Catalog Number:<br>18736-1-AP  | GenBank Accession Number:<br>BC 148863              |                | Purification Method:<br>Antigen affinity purification   |  |
|------------------------|--|---|----------------|---|--|
|                        | Size: Genel D (NCBI):   500 μ g/ml 3239   5 5  |   | BI):           | Recommended Dilutions:<br>WB 1:1000-1:4000<br>IP 0.5-4.0 ug for 1.0-3.0 mg of total<br>protein lysate<br>IHC 1:20-1:200 |  |
|                        |  |   |                |   |  |
|                        | Source:<br>Rabbit  | UNIPROT ID:<br>P35453<br>Full Name:<br>homeobox D13 |                |   |  |
|                        | lsotype:<br>IgG  |   |                |   |  |
|                        |  | Calculated M<br>343 aa, 36 k                        |                |   |  |
|                        |  | Observed MW:<br>36 kDa                              |                |   |  |
| Applications           | Tested Applications:   |   |                | ontrols:  |  |
|                        | Cited Applications: lu   |   |                | : A 549 cells, rat brain tissue, Jurkat cells, mouse<br>g tissue, mouse brain tissue                                    |  |
|                        | WB IP: A54<br>Species Specificity:   |   |                | ells,   |  |
|                        | human mouse rat  |   |                | ıman brain tissue, human hepatocirrhosis<br>human liver cancer tissue   |  |
|                        | Cited Species:<br>human, mouse   |   |                |   |  |
|                        | Note-IHC: suggested antigen retrieval with<br>TE buffer pH 9.0; (*) Alternatively, antigen<br>retrieval may be performed with citrate<br>buffer pH 6.0   |   |                |   |  |
| Background Information | HOXD13, also named as HOX4I, belongs to the Abd-B homeobox family. HOXD13 is a sequence-specific transcription<br>factor which is part of a developmental regulatory system that provides cells with specific positional identities on<br>the anterior-posterior axis. Mutation of HOXD13 will cause synpolydactyly (SPD), brachydactyly type D (BDD),<br>syndactyly type 5brachydactyly-syndactyly syndrome (BDSD) or brachydactyly type E (BDE).Present polyclonal and<br>HOX13D antibody(18736-1-AP) is produced by immunizing animals with part of N-terminus chain of HOX13D and<br>detect a 36-kDa band in cell and tissues. |   |                |   |  |
| Notable Publications   | Author   | Pubmed ID   | Journal        | Application   |  |
|                        | Fei-Yang Jin   | 30431285  | Mol Pharm      |   |  |
|                        | Xingyue Ma   | 39744569  | J Cancer       | WB  |  |
|                        | Lishan Zhang   | 36804539  | Cell Death Dis | WB  |  |
| Storage                | Storage:   | e year after shipment.                              |                |   |  |

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data






A 549 cells were subjected to SDS PAGE followed by western blot with 18736-1-AP (HOXD13 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human brain using 18736-1-AP (HOXD13 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human brain using 18736-1-AP (HOXD13 antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-HOXD13 (IP:18736-1-AP, 4ug; Detection:18736-1-AP 1:500) with A549 cells lysate 1600ug.