For Research Use Only

PSKH2 Polyclonal antibody

Catalog Number:18813-1-AP




| Basic Information | Catalog Number: 18813-1-AP | GenBank Accession Number: BC126182 | | Purification Method: Antigen affinity purification | |
|------------------------|---|---|---|--|--|
| | Size: 400 µg/ml | GenelD (NCBI): 85481 | | Recommended Dilutions: WB 1:1000-1:4000 | |
| | Source: Rabbit | UNIPROT ID: Q96QS6 Full Name: protein serine kinase H2 Calculated MW: 43 kDa | | IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200 IF 1:20-1:200 | |
| | Isotype: IgG | | | | |
| | | | | | |
| | | Observed MW: 35 kDa | | | |
| Applications | Tested Applications: IF/ICC. IHC. IP. WB.ELISA | Tested Applications: IF/ICC, IHC, IP, WB,ELISA | | Positive Controls: | |
| | Species Specificity: | | WB : HepG2 cells, human brain tissue, human heart tissue, human testis tissue | | |
| | human | Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen | | ecells, | |
| | | | | IHC : human kidney tissue, | |
| | | | | 2 cells, | |
| Background Information | J | PSKH2 belongs to the protein kinase superfamily and CAMK Ser/Thr protein kinase family. PSKH2 catalytic the reaction: ATP + a protein = ADP + a phosphoprotein. | | | |
| Storage | Storage: Store at -20°C. Stable for one Storage Buffer: PBS with 0.02% sodium azic Aliquoting is unnecessary fo | le and 50% glycerol pH 7.3. | | | |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 18813-1-AP (PSKH2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human kidney using 18813-1-AP (PSKH2 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney using 18813-1-AP (PSKH2 antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-PSKH2 (IP:18813-1-AP, 4ug; Detection:18813-1-AP 1:500) with HepG2 cells lysate 3400ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HepC2 cells using 18813-1-AP (PSKH2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).