## For Research Use Only

# IRF8 Polyclonal antibody

Catalog Number: 18977-1-AP 17 Publications



#### **Basic Information**

Catalog Number: 18977-1-AP Concentration: 1200 µ g/ml Source: Rabbit Isotype: GenBank Accession Number: NM\_002163 GeneID (NCBI): 3394 UNIPROT ID: Q02556 Full Name: ICSBP1 Calculated MW:

Purification Method: Antigen affinity purification Recommended Dilutions: WB: 1:500-1:2000 IHC: 1:50-1:200

## **Applications**

Tested Applications: WB, IHC, ELISA
Cited Applications: WB, IHC, IF
Species Specificity: human, mouse, rat
Cited Species: human, mouse, rat

editaty: Duse, rat ies:

48 kDa Observed MW: 48 kDa

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

# **Background Information**

IRF8 also named as ICSBP1, is a transcription factor of the IFN regulatory factor (IRF) family. The IRF family proteins bind to the IFN-stimulated response element (ISRE) and regulate expression of genes stimulated by type I IFNs, namely IFN-alpha and IFN-beta. IRF family proteins also control expression of IFN-alpha and IFN-beta-regulated genes that are induced by viral infection. IRF8 specifically binds to the upstream regulatory region of type I IFN and IFN-inducible MHC class I genes (the IFN consensus sequence (ICS)). It plays a negative regulatory role in cells of the immune system. It is predominantly in lymphoid tissues.

Positive Controls:

WB: mouse spleen tissue,

IHC: human lymphoma tissue,

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yide He	31753358	Mater Sci Eng C Mater Biol Appl	WB,IF
Yanhui Jia	36356459	Immunobiology	WB
Xin Wang	35070752	Transl Lung Cancer Res	IHC

# Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


# Selected Validation Data



mouse spleen tissue were subjected to SDS PAGE followed by western blot with 18977-1-AP (IRF8 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lymphoma using 18977-1-AP (IRF8 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human lymphoma using 18977-1-AP (IRF8 antibody) at dilution of 1:100 (under 40x lens).