

For Research Use Only

NOX2 Polyclonal antibody

Catalog Number: 19013-1-AP

Featured Product

150 Publications



Basic Information

Catalog Number: 19013-1-AP	GenBank Accession Number: BC032720	Purification Method: Antigen affinity purification
Size: 650 µg/ml	GeneID (NCBI): 1536	Recommended Dilutions: WB 1:2000-1:8000 IHC 1:500-1:2000 IF 1:50-1:500
Source: Rabbit	UNIPROT ID: P04839	
Isotype: IgG	Full Name: cytochrome b-245, beta polypeptide	
Immunogen Catalog Number: AG5536	Calculated MW: 577 aa, 65 kDa	
	Observed MW: 55 kDa	

Applications

Tested Applications: FC, IF/ICC, IF-P, IHC, WB, ELISA	Positive Controls: WB : mouse spleen tissue, Jurkat cells, Raji cells, human plasma, rat spleen tissue IHC : human tonsillitis tissue, mouse spleen tissue IF : mouse brain tissue, human tonsillitis tissue
Cited Applications: WB, IF, IHC, CoIP	
Species Specificity: human, mouse, rat	
Cited Species: human, rat, mouse, rabbit, pig, canine	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

NOX2, also named as CYBB, CGD, 91-phox, gp91-1, gp91-phox, p22 phagocyte B-cytochrome, cytochrome b-245 and beta polypeptide, is a critical component of the membrane-bound oxidase of phagocytes that generates superoxide. It is the terminal component of a respiratory chain that transfers single electrons from cytoplasmic NADPH across the plasma membrane to molecular oxygen on the exterior. This full length protein has three glycosylation sites. CYBB is found in human cardiomyocytes as multiple bands: the signal between 55 and 65 kDa is probably the unglycosylated protein, because the predicted molecular weight of unglycosylated CYBB protein is approximately 58 kDa (PMID: 17587483) to 65 kDa in phagocytes and the bands around 80 kDa probably represent glycosylated CYBB, as has also been shown in human umbilical vein endothelial cells (PMID:12610097). In other reports, it also can be detected a band of 91 kDa (PMID:19965781).

Notable Publications

Author	Pubmed ID	Journal	Application
Zi-Chao Wang	36163178	Cell Death Dis	WB
Jingyu Song	31564534	J Sex Med	
Jing Li	31616516	Exp Ther Med	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 19013-1-AP (NOX2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




mouse spleen tissue were subjected to SDS PAGE followed by western blot with 19013-1-AP (NOX2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 19013-1-AP (NOX2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 19013-1-AP (NOX2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 19013-1-AP (NOX2 antibody) at dilution of 1:1000 (under 4x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using NOX2 antibody (19013-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10⁶ RAW 264.7 cells were intracellularly stained with 0.4 ug Anti-Human NOX2 (19013-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).