

For Research Use Only

KLC1 Polyclonal antibody, PBS Only

Catalog Number:19028-1-PBS

Featured Product



Basic Information

Catalog Number:

19028-1-PBS

Concentration:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG13570

GenBank Accession Number:

BC008881

GeneID (NCBI):

3831

UNIPROT ID:

Q07866

Full Name:

kinesin light chain 1

Calculated MW:

65 kDa

Observed MW:

61-70 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

Kinesin is a microtubule-associated force-producing protein that may play a role in organelle transport. Kinesin is tetrameric proteins containing two heavy (alpha) chains and two light (beta) chains. The alpha chains provide the tubulin binding site and the ATPase domains, whereas the beta chains are responsible for the specific attachment of the organelle to be moved by the kinesin tetramer (PubMed:21385839).

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 19028-1-AP (KLC1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 19028-1-PBS in a different storage buffer formulation.




mouse brain tissue were subjected to SDS PAGE followed by western blot with 19028-1-AP (KLC1 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 19028-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human Bowen disease slide using 19028-1-AP (KLC1 antibody) at dilution of 1:300 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 19028-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 19028-1-AP (KLC1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 19028-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 19028-1-AP (KLC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 19028-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using KLC1 antibody (19028-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 19028-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using KLC1 antibody (19028-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 19028-1-PBS in a different storage buffer formulation.