For Research Use Only

MDM2 Polyclonal antibody

Catalog Number:19058-1-AP






Basic Information	Catalog Number: 19058-1-AP	GenBank Accession Num NM_002392		urification Method: Intigen affinity purification
	Size:	GeneID (NCBI):	R	ecommended Dilutions:
	300 µg/ml	4193	v	VB 1:200-1:1000
	Source: UNIPROTID: Rabbit 000987			
	Isotype: Full Name:			
	IgG Mdm2 p53 binding protein homolog (mouse)			
		Calculated MW: 55 kDa		
	Observed MW: 30-55 kDa, 60-100 kDa			
Applications	Tested Applications:	Positive Controls:		
	WB, ELISA Cited Applications: WB, IHC, IF, IP, COIP	pplications:		
	Species Specificity: human, mouse, rat			
	Cited Species: human, mouse, rat, artemia sinica			
Background Information	MDM2, also known as HDM2, is a protein of the E3 ubiquitin-ligase family. MDM2 is recognized as a proto-oncogene and is over-expressed in a wide range of human malignancies including soft tissue carcinomas and breast cancer. MDM2 can bind to the p53 tumor suppressor protein with high affinity, and negatively regulates p53 by mediating its ubiquitination and proteosomal degradation (PMID: 17000718). MDM2 has some isoforms with calculated molecular mass of 25-55kD, and can be detected as 90-97kD. In addition to full length MDM2, two lower bands (85kl and 75kD) are often observed which are likely to be produced by translation initiation at methionine codons 50 and 60. Another isoform of MDM2 is often observed as 57-60kD in cell lines which is indeed a caspase cleavage product of MDM2 at residue 361 (PMID: 19565011, 9840926, 7689721).			
Notable Publications	Author P	ubmed ID Journal		Application
		1839745 Int J Med	l Sci	WB
		1626714 FEBS Let		WB,IP
		4650049 Cell Dea		WB

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



A431 cells were subjected to SDS PAGE followed by western blot with 19058-1-AP (MDM2 Antibody) at dilution of 1:300 incubated at 4 degree celsius over night.