

For Research Use Only

FOXO4 Polyclonal antibody

Catalog Number: 19121-1-AP

1 Publications



Basic Information

Catalog Number:

19121-1-AP

Size:

230 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_005938

GeneID (NCBI):

4303

UNIPROT ID:

P98177

Full Name:

forkhead box O4

Calculated MW:

54 kDa

Observed MW:

45-48 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IF/ICC 1:10-1:100

Applications

Tested Applications:

IF/ICC, WB, ELISA

Cited Applications:

WB, IF

Species Specificity:

human, mouse

Cited Species:

human, mouse

Positive Controls:

WB : human heart tissue, mouse heart tissue

IF/ICC : HepG2 cells,

Background Information

FOXO4, also named as AFX, AFX1 and MLLT7, is a transcription factor which involved in the regulation of the ins signaling pathway. It binds to ins-response elements (IREs) and can activate transcription of IGFBP1. FOXO down-regulates expression of HIF1A and suppresses hypoxia-induced transcriptional activation of HIF1A-modulated genes. It is involved in negative regulation of the cell cycle. A chromosomal aberration involving FOXO4 is found in acute leukemias. The antibody can recognize isoform 1 and isoform Zeta of FOXO4.

Notable Publications

Author	Pubmed ID	Journal	Application
Xi-Lin Liu	33838260	Neurosci Lett	WB, IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human heart tissue were subjected to SDS PAGE followed by western blot with 19121-1-AP (FOXO4 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using FOXO4 antibody 19121-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).