For Research Use Only

DCC-Specific Polyclonal antibody

Catalog Number:19123-1-AP

3 Publications




Basic Information	Catalog Number: 19123-1-AP	GenBank Accession NM_005215		ication Method: gen affinity purification	
	Size:	GenelD (NCBI):		mmended Dilutions:	
	850 µg/ml	5		:500-1:1000	
	Source: Rabbit	UNIPROT ID: P43146			
	Isotype: Full Name: IgG deleted in colorectal carcinoma Calculated MW: 1447 aa, 158 kDa				
	Observed MW: 180 kDa				
Applications	Tested Applications: WB,ELISA		Positive Controls:		
	WBLEISA WB : mouse brain tissue, Cited Applications: WB, IF, IHC				
	Species Specificity: human, mouse, rat				
	Cited Species: rat, mouse				
Background Information	DCC, also named as IGDCC1, belongs to the immunoglobulin superfamily and DCC family. DCC is a receptor for netrin required for axon guidance. DCC mediates axon attraction of neuronal growth cones in the developing nervous system upon ligand binding. Its association with UNC5 proteins may trigger signaling for axon repulsion. I also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand. DCC gene is a tumor suppressor gene. This antibody raised against a synthesized polypeptide of human DCC protein recognizes endogenous DCC with an apparent molecular weight of 180 kDa. As the predicted molecular mass of DCC is 158 kDa, the slower mobility is likely attributable to post-translational modification, perhaps by glycosylation at potential N-linked glycosylation sites (PMID: 7926722; 8044801).				
Notable Publications	Author	Pubmed ID Jo	ournal	Application	
	Mian Xiao	34344989 S	ci Rep	IF	
	Dongdong Chai	32271715 A	ging (Albany NY)	WB	
	Songjie Ji	38628640 F	ront Pharmacol	WB,IHC	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 19123-1-AP (DCC-Specific Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.