

For Research Use Only

ATG14/Barkor (N-terminal) Polyclonal antibody

Catalog Number: 19491-1-AP

Featured Product

34 Publications



Basic Information

Catalog Number:

19491-1-AP

Concentration:

350 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_014924

GeneID (NCBI):

22863

UNIPROT ID:

Q6ZNE5

Full Name:

KIAA0831

Calculated MW:

55 kDa

Observed MW:

65-68 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IF, IP

Species Specificity:

human, mouse

Cited Species:

human, mouse, rat, pig, canine, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HEK-293 cells, A431 cells, mouse brain tissue, NIH/3T3 cells

IHC : mouse brain tissue,

IF/ICC : MCF-7 cells,

Background Information

Barkor, also named as KIAA0831, Atg14 and Atg14L, is required for both basal and inducible autophagy. It plays a role in autophagosome formation and MAP1LC3/LC3 conjugation to phosphatidylethanolamine. Barkor promotes BECN1 translocation from the trans-Golgi network to autophagosomes. This antibody is specific to KIAA0831(Barkor).

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------------|-----------|-----------------------------------|-------------|
| Luis Muniz-Feliciano | 28933590 | Autophagy | WB |
| Qiang Fu | 25234643 | J Gen Virol | WB |
| Xueqiang Peng | 32949647 | Biochim Biophys Acta Mol Cell Res | WB |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 19491-1-AP (ATG14/Barkor (N-terminal) antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 19491-1-AP (ATG14/Barkor (N-terminal) antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using 19491-1-AP (ATG14/Barkor (N-terminal) antibody) at dilution of 1:50 and CoraLite488-Conjugated Goat Anti-Rabbit IgG(H+L).