For Research Use Only

TRAPPC9/NIBP Polyclonal antibody

Catalog Number: 19549-1-AP




Basic Information	Catalog Number: 19549-1-AP	GenBank Accession Number: NM_031466	Purification Method: Antigen affinity purification		
	Size: 170 µ g/ml Source: Rabbit Isotype: IgG	GeneID (NCBI): 83696 UNIPROT ID: Q96Q05	Recommended Dilutions: WB 1:500-1:3000 IHC 1:20-1:200		
				Full Name: trafficking protein particle complex 9 Calculated MW: 139 kDa	
		Observed MW: 56 kDa			
		Applications	Tested Applications: IHC, WB,ELISA Species Specificity: human, mouse, rat		ve Controls:
					WB : SH-SY5Y cells, human brain tissue, mouse kidney tissue
IHC : h	uman heart tissue, human kidney tissue				
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	TRAPPC9, also named NIK- and IKBKB-binding protein (NIBP), is involved in the NF-kappaB signaling pathway and directly interacts with IKK-beta and MAP3K14. TRAPPC9 functions as an activator of NF-kappa-B via increased phosphorylation of the IKK complex. TARPPC9 may also function in neuronal cells differentiation and defects in TRAPPC9 are the cause of mental retardation autosomal recessive type 13 (MRT13), which is characterized by significantly below average general intellectual functioning associated with impairments in adaptative behavior. TARPPC9 is a component of the multisubunit TRAPP (transport protein particle) complex and it may play a role in vesicular transport from endoplasmic reticulum to Golgi.				
Storage	Storage: Store at -20°C. Stable for one Storage Buffer: PBS with 0.02% sodium azide Aliquoting is unnecessary for	and 50% glycerol pH 7.3.			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 19549-1-AP (TRAPPC9/NIBP antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human heart using 19549-1-AP (TRAPPC9/NIBP antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human heart using 19549-1-AP (TRAPPC9/NIBP antibody) at dilution of 1:100 (under 40x lens).