For Research Use Only

SATB2 Polyclonal antibody

Catalog Number: 19557-1-AP



Basic Information

Catalog Number: 19557-1-AP

Isotype:

NM_015265 GeneID (NCBI):

GenBank Accession Number:

Size: 250 ug/ml Source: Rabbit

23314 UNIPROT ID: Q9UPW6 Full Name:

Calculated MW: 83 kDa Observed MW: ~100 kDa

SATB homeobox 2

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

IHC 1:50-1:500

Applications

Tested Applications: WB, IHC, ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human, mouse

Positive Controls:

WB: HepG2 cells,

IHC: human colon cancer tissue,

Background Information

SATB2, also named as KIAA1034, belongs to the CUT homeobox family. SATB2 binds to DNA at nuclear matrix- or scaffold-associated regions. STAB2 recognizes the sugar-phosphate structure of double-stranded DNA. SATB2 is a transcription factor controlling nuclear gene expression, by binding to matrix attachment regions (MARs) of DNA and inducing a local chromatin-loop remodeling. SATB2 acts as a docking site for several chromatin remodeling enzymes and also by recruiting corepressors (HDACs) or coactivators (HATs) directly to promoters and enhancers. It is required for the initiation of the upper-layer neurons (UL1) specific genetic program and for the inactivation of deep-layer neurons (DL) and UL2 specific genes, probably by modulating BCL11B expression. It is a repressor of Ctip2 and regulatory determinant of corticocortical connections in the developing cerebral cortex. SATB2 may play an important role in palate formation. SATB2 acts as a molecular node in a transcriptional network regulating skeletal development and osteoblast differentiation.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



The lysate of total HeLa cells and nuclear extracted from HeLa cells were subjected to SDS PAGE followed by western blot with 19557-1-AP (SATB2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded Colorectal cancer slide using 19557-1-AP (SATB2 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).