## For Research Use Only

# **UNC45A Polyclonal antibody**

Catalog Number: 19564-1-AP

Featured Product

**3 Publications** 



**Basic Information** 

Catalog Number: GenBank Accession Number: 19564-1-AP NM\_018671
Size: GeneID (NCBI): 600 µ g/ml 55898
Source: UNIPROT ID: Rabbit Q9H3U1

Rabbit Q9H3U1 Isotype: Full Name:

gG unc-45 homolog A (C. elegans)

Calculated MW: 103 kDa Observed MW: 100 kDa Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

**Applications** 

Tested Applications: IHC, IP, WB,ELISA Cited Applications: WB, IF, IHC Species Specificity:

human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: K-562 cells, mouse lung tissue, mouse ovary tissue. HeLa cells

IP: K-562 cells,

IHC: human ovary tumor tissue, human tonsil tissue

# **Background Information**

UNC45A, also named as SMAP1, acts as co-chaperone for HSP90. UNC45A prevents the stimulation of HSP90AB1 ATPase activity by AHSA1. It is a positive factor in promoting PGR function in the cell. It is necessary for proper folding of myosin and necessary for normal cell proliferation, normal myotube formation and myosin accumulation during muscle cell development. UNC45A plays a role in erythropoiesis in stroma cells in the spleen. The antibody recognizes the C-term of UNC45A.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Susana Lechuga	35344227	FASEB J	WB
Valentino Clemente	34206743	Cells	IHC,IF
Juri Habicht	33262310	J Cell Sci	WB, IF

# Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


# **Selected Validation Data**




K-562 cells were subjected to SDS PAGE followed by western blot with 19564-1-AP (UNC 45A antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



WB result of UNC45A antibody (19564-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-UNC45A transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 19564-1-AP (UNC 45A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-UNC 45A (IP:19564-1-AP, 4ug; Detection:19564-1-AP 1:300) with K-562 cells lysate 3600ug.