For Research Use Only

KCC2/SLC12A5-Specific Polyclonal antibody



Catalog Number: 19565-1-AP

6 Publications

Basic Information

Catalog Number:
19565-1-AP
Size:
550 µ g/ml
Source:
Rabbit
Isotype:
IgG

GenBank Accession Number: NM_020708 GeneID (NCBI): 57468

UNIPROT ID: Q9H2X9 Full Name:

solute carrier family 12 (potassiumchloride transporter), member 5

Calculated MW: 126 kDa Observed MW: 130-150 kDa

Applications

Tested Applications: IF-P, IHC, WB, ELISA Cited Applications: WB, IF, IHC Species Specificity: human, mouse, rat Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human brain tissue, mouse brain tissue, rat brain tissue

Purification Method:

WB 1:500-1:2000 IHC 1:20-1:200

IF-P 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC : human brain tissue, rat brain tissue, mouse brain

tissue

IF-P: mouse cerebellum tissue,

Background Information

KCC2, encoded by SLC12A5, is a neuron-specific K+-Cl- cotransporter that is essential for Cl- homeostasis and inhibitory synaptic transmission in brain. KCC2 is expressed at high levels in neurons throughout the nervous system. KCC2 deletion caused animal death. Alterations of KCC2 have been linked to neuropathic pain, temporal lobe epilepsy, and other diseases. This antibody can detect 130-150 kDa monomer and ~250 kDa dimer of KKC2 in brain lysate. (PMID: 23401525)

Notable Publications

Author	Pubmed ID	Journal	Application
Aaron J Barbour	34445881	ASN Neuro	WB, IF
Shuo Zhao	35106976	Brain Behav	WB
Aaron J Barbour	32344154	Neurobiol Dis	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.cor

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




human brain tissue were subjected to SDS PAGE followed by western blot with 19565-1-AP (KCC2/SLC12A5-Specific antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 19565-1-AP (KCC2/SLC12A5-Specific antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human brain using 19565-1-AP (KCC2/SLC12A5-Specific antibody) at dilution of 1:100 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human brain using 19565-1-AP (KCC2/SLC12A5-Specific antibody) at dilution of 1:100 (under 40x lens).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 19565-1-AP (KCC2/SLC12A5-Specific antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 19565-1-AP (KCC2/SLC12A5-Specific antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 19565-1-AP (KCC2/SLC12A5-Specific antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse cerebellum tissue using KCC2/SLC12A5-Specific antibody (19565-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse cerebellum tissue using KCC2/SLC12A5-Specific antibody (19565-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 19565-1-AP (KCC2/SLC12A5-Specific antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).