For Research Use Only

ND1 Polyclonal antibody Catalog Number: 19703-1-AP Featured Product

Featured Product







Basic Information	Catalog Number: 19703-1-AP	GenBank Accession Number: NC_012920		Purification Method: Antigen affinity purification	
	Concentration:GeneID (NCBI):600 ug/ml4535Source:UNIPROT ID:RabbitP03886			Recommended Dilutions: WB 1:1000-1:8000	
			IHC 1:50-1:500		
	Isotype: Full Name: IgG NADH dehydrogenase, subunit 1 (complex I)				
		Calculated MW: 36 kDa			
	Observed MW: 28-38 kDa				
Applications	Tested Applications: WB, IHC, ELISA		Positive Controls:		
	Cited Applications: WB, IHC, IF	ted Applications: heart tissue, mouse kidney tissue, mouse liver tissue,		mouse kidney tissue, mouse liver	
	Species Specificity: IHC : human liver tissue, human, mouse IHC : human liver tissue,				
	Cited Species: human, mouse, rat, pig, hamster				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	ND1 belongs to the complex I subunit 1 family. ND1 is a core subunit of the mitochondrial membrane respiraton chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for cataly Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.				
Notable Publications	Author	Pubmed ID Journal Application			
	Q E Xie	34571250 Mi	tochondrion	WB	
	Shibo Bao	36116352 Ec	otoxicol Environ S	af WB	
	Christopher Lowden	34525369 Ce	ell Rep	WB	
Storage	Storage: Store at -20°C. Stable for on Storage Buffer: PBS with 0.02% sodium azi	e year after shipment. de and 50% glycerol pH 7.3.			

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 19703-1-AP (ND1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Various lysates were subjected to SDS PAGE followed by western blot with 19703-1-AP (ND1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 19703-1-AP (ND1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).