For Research Use Only

PYGM-Specific Polyclonal antibody

Catalog Number: 19716-1-AP

Featured Product

10 Publications

NM_005609

UNIPROT ID:

5837

P11217 Full Name:

GeneID (NCBI):

GenBank Accession Number:

phosphorylase, glycogen, muscle



Basic Information

Catalog Number: 19716-1-AP Size:

600 ug/ml Source: Rabbit Isotype:

Calculated MW: 97 kDa

> Observed MW: 97-105 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:2000-1:10000

IHC 1:20-1:200

IF/ICC 1:200-1:800

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC

Species Specificity: human, mouse, rat Cited Species: human, mouse, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse heart tissue, rat heart

IHC: human skeletal muscle tissue, human liver tissue

IF/ICC: HepG2 cells,

Background Information

PYGM belongs to the glycogen phosphorylase family. Phosphorylase is an important allosteric enzyme in carbohydrate metabolism. Enzymes from different sources differ in their regulatory mechanisms and in their natural substrates. However, all known phosphorylases share catalytic and structural properties. Defects in PYGM are the cause of glycogen storage disease type 5 (GSD5) which also known as McArdle disease. The antibody is specific to PYGM.

Notable Publications

Author	Pubmed ID	Journal	Application
Omer Keinan	34707293	Nature	WB
Jia-Hua Qu	36359893	Cells	WB
Lingyu Jiang	35761940	Oncol Lett	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 19716-1-AP (PYGM-Specific antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




WB result of PYGM-Specific antibody (19716-1-AP; 1:1200; incubated at room temperature for 1.5 hours) with sh-Control and sh-PYGM-Specific transfected HepG2 cells.



Immunohistochemical analysis of paraffinembedded human skeletal muscle using 19716-1-AP (PYGM-Specific antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human skeletal muscle using 19716-1-AP (PYGM-Specific antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using PYGM-Specific antibody (19716-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).