## For Research Use Only FUNDC2 Polyclonal antibody Catalog Number: 19832-1-AP



Basic Information	Catalog Number: 19832-1-AP	GenBank Accession Number: BC000255	Purification Method: Antigen affinity purification	
	Size: 400 ug/ml Source: Rabbit	GeneID (NCBI): 65991 UNIPROT ID: Q9BWH2	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500	
				lsotype: IgG
	Immunogen Catalog Number: AG13800	Calculated MW: 189 aa, 21 kDa		
		Observed MW: 21 kDa		
	Applications	Tested Applications: WB, IHC, ELISA	Positive Controls:	
Species Specificity: human		we : HeLa cells, HuH-7 cells, HePu2 cells becies Specificity:		
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Alterno retrieval may be performed buffer pH 6.0	atively, antigen		
Background Information	<ul> <li>TE buffer pH 9.0; (*) Alternor retrieval may be performed buffer pH 6.0</li> <li>FUNDC2, or FUN14 domain-contai role in various cellular processes, implicated in the progression of se liver cancer. In the context of cance prognosis in hepatocellular carcin mediated mitochondrial fusion, w plays a role in platelet function ar which is a key lipid second messe</li> </ul>	ning protein 2, is a mitochondrial oute including the regulation of cell surviv everal types of cancer, particularly in i ere, FUNDC2 has been shown to promo oma (HCC) and is involved in mitocho hich can lead to changes in cellular m d survival. It binds directly to phosph nger in cell signaling, and this interacc has been shown to decrease platelet	er membrane protein that plays a significan al, apoptosis, and metabolism. It has been triple-negative breast cancer (TNBC) and te tumorigenesis. It is associated with poor ondrial fragmentation by inhibiting MFN1- etabolism and energy levels. FUNDC2 also atidylinositol-3,4,5-trisphosphate (PIP3), tion is crucial for the recruitment of PIP3 to aggregation and impair hemostasis in mice	

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 19832-1-AP (FUNDC2 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 19832-1-AP (FUNDC2 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Various lysates were subjected to SDS PAGE followed by western blot with 19832-1-AP (FUNDC2 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.