For Research Use Only

KTN1 Polyclonal antibody

Catalog Number: 19841-1-AP

Featured Product

8 Publications

BC117132

GenBank Accession Number:



Basic Information

Catalog Number: 19841-1-AP

Size: GeneID (NCBI): 1000 μ g/ml 3895

Source: UNIPROT ID:
Rabbit Q86UP2
Isotype: Full Name:

gG kinectin 1 (kinesin receptor)

Immunogen Catalog Number: Calculated MW: AG13854 1357 aa, 156 kDa

Observed MW: 156 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF 1:200-1:800

Applications

Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

Cited Applications: CoIP, IF, IHC, WB Species Specificity: human, mouse Cited Species:

human, mouse, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, A549 cells, Jurkat cells, K-562 cells,

PC-3 cells, SKOV-3 cells

IP: Jurkat cells,

IHC: human ovary tumor tissue, human testis tissue

IF: HepG2 cells,

Background Information

KTN1/Kinectin is an integral endoplasmic reticulum (ER) membrane protein which acts as dynamic anchor to facilitate ER membrane extension along microtubules. It has been found to bind kinesin, Rho GTPase, and translation elongation factor-1delta and gets involved in organelle motility and protein synthesis.

Notable Publications

Author	Pubmed ID	Journal	Application
Sara Picó	34586830	Sci Transl Med	WB
Pengli Zheng	30030520	Cell Res	IF
Qingzhou Chen	30773365	Curr Biol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 19841-1-AP (KTN1 antibody) at dilution of 1:8000 incubated at room temporarium for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 19841-1-AP (KTN1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-KTN1 (IP:19841-1-AP, 4ug; Detection:19841-1-AP 1:500) with Jurkat cells lysate 1200ug.



WB result of KTN1 antibody (19841-1-AP; 1:12000; incubated at room temperature for 1.5 hours) with sh-Control and sh-KTN1 transfected A549 cells.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using KTN1 antibody (19841-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using KTN1 antibody (19841-1-AP) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).