

For Research Use Only

NF90/ILF3 Polyclonal antibody

Catalog Number: 19887-1-AP

Featured Product

29 Publications



Basic Information

Catalog Number:

19887-1-AP

Size:

700 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG13707

GenBank Accession Number:

BC064836

GeneID (NCBI):

3609

UNIPROT ID:

Q12906

Full Name:

interleukin enhancer binding factor 3,
90kDa

Calculated MW:

894 aa, 95 kDa

Observed MW:

90 kDa, 110 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total
protein lysate

IHC 1:20-1:200

IF/ICC 1:10-1:100

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, RIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : HeLa cells, Jurkat cells, mouse spleen tissue, K-
562 cells, Raji cells

IP : HeLa cells,

IHC : human cervical cancer tissue, human placenta
tissue

IF/ICC : HeLa cells,

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Background Information

Nuclear factor of activated T cells (NFAT) is a transcription factor required for T-cell expression of interleukin-2 (IL2). NFAT binds to a sequence known as the antigen receptor response element-2 (ARRE2) in the IL2 enhancer. NFAT contains two subunits, one is ILF3. ILF3 can facilitate double-stranded RNA-regulated gene expression at the level of post-transcription. It binds to and inhibits coding sequences of acid beta-glucosidase and other mRNAs and functions at the initiation phase of GCaSe mRNA translation, probably by inhibiting its binding to polysomes. ILF3 exists in two isoforms NFAR-1 (90 kDa) and -2 (110 kDa). NFAR-1 and -2 each contain two DRBDs in their C-terminus regions and exhibit 98% homology to one another at the amino acid level. NFAR-2 contains an extra 192 amino acids at the C-terminal region that are encoded by an extra three exons present in its corresponding transcript.

Notable Publications

Author	Pubmed ID	Journal	Application
Ling-Fei Zhang	26346275	EMBO J	WB
Neeraja Purandare	36339251	iScience	WB
Subhayan Sur	33043604	EMBO Rep	WB, RIP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 19887-1-AP (ILF3 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



IP result of anti-NF90/ILF3 (IP:19887-1-AP, 3ug; Detection:19887-1-AP 1:1000) with HeLa cells lysate 1600ug.



Immunohistochemical analysis of paraffin-embedded human cervical cancer using 19887-1-AP (ILF3 antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of HeLa cells, using ILF3 antibody 19887-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).