### For Research Use Only

# GLI3-Specific Polyclonal antibody

Catalog Number: 19949-1-AP

12 Publications



**Basic Information** 

Catalog Number: 19949-1-AP Size:

240 µg/ml Source: Rabbit Isotype:

IgG

GenBank Accession Number:

NM\_000168
GeneID (NCBI):
2737
UNIPROT ID:
P10071
Full Name:

GLI family zinc finger 3
Calculated MW:

Calculated MW: 170 kDa Observed MW: 190 kDa, 83-86 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:200-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200 IF/ICC 1:10-1:100

**Applications** 

Tested Applications:

WB, IP, IF/ICC, IHC, ELISA

Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: human placenta tissue, mouse lung tissue

IP: mouse lung tissue,

IHC: human testis tissue, human colon tissue

IF/ICC: HepG2 cells,

## **Background Information**

GLI3 belongs to the GLI C2H2-type zinc-finger protein family. GLI3 plays a role in limb and brain development. GLI3 is implicated in the transduction of SHH signal. Defects in GLI3 are the cause of Greig cephalo-poly-syndactyly syndrome (GCPS). Defects in GLI3 are a cause of Pallister-Hall syndrome (PHS). Defects in GLI3 are a cause of type A1/B postaxial polydactyly (PAPA1/PAPB). Defects in GLI3 are a cause of type IV preaxial polydactyly. Defects in GLI3 are the cause of acrocallosal syndrome (ACS). The antibody is specific to GLI3. At the molecular level, Gli3 is translated into a 190-kDa transcriptional activator (Gli3-190) that undergoes proteolytic processing into a truncated 83-kDa repressor (Gli3-83) lacking C-terminal activation domains. (PMID: 16705181)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Diana Trnski	26385428	Biochim Biophys Acta	WB
Yuqin Men	26549569	Sci Rep	WB
Petar Ozretić	29039491	Int J Oncol	WB

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.cor

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




human placenta tissue were subjected to SDS PAGE followed by western blot with 19949-1-AP (GLI3-Specific antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells using 19949-1-AP (GLI3-Specific antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.



IP result of anti-GLI3-Specific (IP:19949-1-AP, 4ug; Detection:19949-1-AP 1:300) with mouse lung tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 19949-1-AP (GLI3-Specific Antibody) at dilution of 1:50.