For Research Use Only

GNRHR Polyclonal antibody

Catalog Number: 19950-1-AP 27 Publications



Basic Information

Catalog Number:
19950-1-AP
Concentration:
800 µg/ml
Source:
Rabbit
Isotype:

GenBank Accession Number: NM_000406 GeneID (NCBI): 2798 UNIPROT ID: P30968 Full Name: GnRH receptor Calculated MW:

38 kDa Observed MW: 60-70 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:6000 IHC 1:500-1:2000

Applications

Tested Applications: WB, IHC, FC (Intra), ELISA Cited Applications: WB, IHC, IF

Species Specificity: human, mouse Cited Species:

human, mouse, rat, pig, sheep

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: MCF-7 cells, mouse ovary tissue

IHC: mouse ovary tissue, human testis tissue, human ovary tissue, mouse testis tissue

Background Information

GNRHR, also named GRHR, belongs to the G-protein coupled receptor 1 family. GNRHR is a receptor for GnRH that mediates the action of GnRH to stimulate the secretion of the gonadotropic hormones luteinizing hormone (LH) and follicle-stimulating hormone (FSH). It mediates its action by association with G-proteins that activate a phosphatidylinositol-calcium second messenger system. Isoform2 of GNRHR may act as an inhibitor of GnRH-R signaling. Defects in GNRHR are a cause of idiopathic hypogonadotropic hypogonadism (IHH). Defects in GNRHR are a cause of fertile eunuch syndrome. The antibody only recognizes the isoform1 of GNRHR. The predicted unmodified molecular weight of the human GNRHR is -38 kDa, the larger band (50-65 kDa) is likely to represent a glycosylated form of GNRHR.

Notable Publications

Author	Pubmed ID	Journal	Application
Eszter Lajkó	30344773	Beilstein J Org Chem	
Ivan Ranđelović	31557968	Int J Mol Sci	WB,FC
Jingyuan Xiong	36106654	Mol Nutr Food Res	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.coi

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




1X10^6 MCF-7 cells were intracellularly stained with 0.2 ug Anti-Human GNRHR (19950-1-AP) (red) or 0.2 ug Control Antibody (blue), and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Various lysates were subjected to SDS PAGE followed by western blot with 19950-1-AP (GNRHR antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse ovary tissue slide using 19950-1-AP (GNRHR antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse ovary tissue slide using 19950-1-AP (GNRHR antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).