For Research Use Only

Kir2.1 Polyclonal antibody

Catalog Number: 19965-1-AP 11 Publications



Basic Information

Catalog Number: 19965-1-AP Concentration: 550 ug/ml

Source: Rabbit Isotype:

IgG

GenBank Accession Number:

NM_000891 GeneID (NCBI): 3759 UNIPROT ID: P63252

Full Name:

potassium inwardly-rectifying channel, subfamily J, member 2

Calculated MW: 48 kDa Observed MW: 50 kDa, 60 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:200-1:1000

Applications

Tested Applications: WB, ELISA

Cited Applications:

WB, IHC

Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, rabbit

Positive Controls:

WB: A549 cells,

Background Information

KCNJ2, also named as HHBIRK1, HHIRK1, IRK1, KIR2.1, LQT7 and SQT3, belongs to the inward rectifier-type potassium channel family. KCNJ2 probably participates in establishing action potential waveform and excitability of neuronal and muscle tissues. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. KCNJ2 can be blocked by extracellular barium or cesium. Defects in KCNJ2 are the cause of long QT syndrome type 7 (LQT7). Defects in KCNJ2 are the cause of short QT syndrome type 3 (SQT3). The antibody recognizes the C-term of KCNJ2.

Notable Publications

Author	Pubmed ID	Journal	Application
Juanjuan Du	32954646	J Cell Mol Med	WB
Zhan Li	28546098	J Mol Cell Cardiol	WB
Weiwei Yu	35729093	Nat Commun	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



A549 cells were subjected to SDS PAGE followed by western blot with 19965-1-AP (Kir2.1 antibody) at dilution of 1:200 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 19965-1-AP (Kir2.1 antibody at dilution of 1:50.