For Research Use Only

Kir2.1 Polyclonal antibody

Catalog Number: 19965-1-AP 13 Publications



Purification Method:

WB: 1:200-1:1000

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number:

19965-1-AP

Source:

Rabbit Isotype:

IgG

GenBank Accession Number:

NM_000891

GeneID (NCBI):

3759

UNIPROT ID: P63252

Full Name:

potassium inwardly-rectifying channel, subfamily J, member 2

Calculated MW:

48 kDa

Observed MW:

50 kDa, 60 kDa

Positive Controls:

WB: A549 cells,

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, rabbit

Background Information

KCNJ2, also named as HHBIRK1, HHIRK1, IRK1, KIR2.1, LQT7 and SQT3, belongs to the inward rectifier-type potassium channel family. KCNJ2 probably participates in establishing action potential waveform and excitability of neuronal and muscle tissues. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. KCNJ2 can be blocked by extracellular barium or cesium. Defects in KCNJ2 are the cause of long QT syndrome type 7 (LQT7). Defects in KCNJ2 are the cause of short QT syndrome type 3 (SQT3). The antibody recognizes the C-term of KCNJ2.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|-------------|-----------|--------------------|-------------|
| Juanjuan Du | 32954646 | J Cell Mol Med | WB |
| Zhan Li | 28546098 | J Mol Cell Cardiol | WB |
| Weiwei Yu | 35729093 | Nat Commun | WB |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



A549 cells were subjected to SDS PAGE followed by western blot with 19965-1-AP (Kir2.1 antibody) at dilution of 1:200 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 19965-1-AP (Kir2.1 antibody at dilution of 1:50.