For Research Use Only

LDHC-Specific Polyclonal antibody

Catalog Number:19989-1-AP 5 Publications



Basic Information

Catalog Number: 19989-1-AP Source:

Isotype: IgG

Rabbit

GenBank Accession Number:

NM_002301 GeneID (NCBI): 3948 UNIPROT ID: P07864 Full Name:

 $lactate\ dehydrogenase\ C$

Calculated MW: 36 kDa Observed MW: 30-36 kDa Purification Method:

Antigen affinity purification Recommended Dilutions:

WB: 1:200-1:1000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:200-1:800 IF/ICC: 1:10-1:100

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC, IF Species Specificity: human, mouse, rat Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

mouse, pig

Positive Controls:

WB: human testis tissue, mouse brain tissue, mouse lung tissue, mouse testis tissue, A549 cells, rat testis tissue

IP: mouse testis tissue,

IHC: mouse testis tissue, human testis tissue, human

heart tissue

IF/ICC: Hela cells,

Background Information

LDHC, also named as CT32 and LDH-X, belongs to the LDH/MDH superfamily and LDH family. LDHC is a possible role in sperm motility. LDHC catalyzes the reaction: (S)-lactate + NAD+ = pyruvate + NADH. LDHC is a testis-specific isoform. Abnormal LDHC expression is associated with several forms of cancer. The antibody has no cross reaction with LDHA and LDHB.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuping Chen	34262451	Front Pharmacol	WB
Li Yang	31332157	Med Sci Monit	IHC
Rie Iida-Norita	36464740	Andrology	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




human testis tissue were subjected to SDS PAGE followed by western blot with 19989-1-AP (LDHC-Specific antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 19989-1-AP (LDHC-Specific antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 19989-1-AP (LDHC-Specific antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of Hela cells, using LDHC antibody 19989-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-LDHC-Specific (IP:19989-1-AP, 4ug; Detection:19989-1-AP 1:500) with mouse testis tissue lysate 2400ug.



Various lysates were subjected to SDS PAGE followed by western blot with 19989-1-AP (LDHC-Specific antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.