

For Research Use Only

CD134/OX40 Polyclonal antibody, PBS Only

Catalog Number: 20006-1-PBS



Basic Information

Catalog Number:	20006-1-PBS	GenBank Accession Number:	NM_003327	Purification Method:
Source:	Rabbit	GenID (NCBI):	7293	Antigen affinity purification
Isotype:	IgG	UNIPROT ID:	P43489	
		Full Name:	tumor necrosis factor receptor superfamily, member 4	
		Calculated MW:	29 kDa	
		Observed MW:	50-55 kDa, 40 kDa	

Applications

Tested Applications:
WB, FC, Indirect ELISA

Species Specificity:
human

Background Information

CD134 (OX40) is a member of the TNFR-superfamily of receptors. Predominantly expressed on activated T cells, CD134 is activated by its cognate ligand CD134L (OX40L) and functions as a T cell co-stimulatory molecule. CD134-CD134L interactions have been proposed as a potential therapeutic target for treating autoimmunity (PMID: 26215166). CD134 can interact with TRAF2, TRAF3, and TRAF5.

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C
Storage Buffer:
PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20006-1-AP (CD134/OX40 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 20006-1-PBS in a different storage buffer formulation.



1X10⁶ day 3 PHA treated human PBMCs were surface stained with 0.2 ug Anti-Human CD134/OX40 (20006-1-AP) and CoraLite®750-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or unstained (blue). Unstimulated human PBMCs were surface stained with 0.2ug Anti-Human CD134/OX40 (20006-1-AP) and CoraLite®750-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (black dashed). Cells were not fixed. Lymphocytes