For Research Use Only

TNS3 Polyclonal antibody

Catalog Number: 20053-1-AP 1 Publications



Basic Information

Catalog Number: 20053-1-AP

Size: 450 μg/ml Source: Rabbit Isotype:

GenBank Accession Number:

NM_022748 GeneID (NCBI): 64759 **UNIPROT ID:** Q68CZ2 Full Name: tensin 3 Calculated MW: 155 kDa

Observed MW: 129 kDa

Purification Method:

IHC 1:50-1:500

Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000

Applications

Tested Applications:

WB, IHC, ELISA **Cited Applications:**

IgG

Species Specificity: human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval antigen retrieval antigen retrieval and 6.0 with citrate buffer pH 6.0

Positive Controls:

WB: mouse kidney tissue, IHC: mouse embryo tissue,

Background Information

TNS3, also named as TEM6, TENS1 and TPP, may play a role in actin remodeling. It is involved in the dissociation of the integrin-tensin-actin complex. EGF activates TNS4 and down-regulates TNS3 which results in capping the tail of ITGB1. TNS3 seems to be involved in mammary cell migration. It may be involved in cell migration and bone development. The antibody recognizes all the isoforms of TNS3.

Notable Publications

Author	Pubmed ID	Journal	Application
Xun Zhang	39382613	Cell Biol Toxicol	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunohistochemical analysis of paraffinembedded mouse embryo tissue slide using 20053-1-AP (TNS3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse embryo tissue slide using 20053-1-AP (TNS3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 20053-1-AP (TNS3 antibody) at dilution of 1:100 incubated at room temperature for 1.5 hours.