For Research Use Only

RDM1 Polyclonal antibody

Catalog Number:20156-1-AP

Featured Product

6 Publications

BC038301

201299

Q8NG50 Full Name:

GeneID (NCBI):

UNIPROT ID:

RAD52 motif 1
Calculated MW:

284 aa, 32 kDa

GenBank Accession Number:



Basic Information

Catalog Number: 20156-1-AP

Size: 340 µ g/ml Source: Rabbit Isotype: IgG

Immunogen Catalog Number: AG14077

G14077

Observed MW: 32 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500 IF 1:20-1:200

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications: WB, IF, IHC

Species Specificity: human

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: PC-3 cells, human placenta tissue, MCF-7 cells

IHC: human lung tissue, human brain tissue, human

testis tissue IF : PC-3 cells,

Background Information

RDM1, also named as RAD52B, encodes a RNA recognition motif (RRM)-containing protein confered resistance to the antitumor agent cisplatin. The expression of RDM1 is closely related to the cell proliferation and apoptosis of thyroid carcinoma cells and RDM1 may be a potential marker of lung adenocarcinoma. RDM1 has many isoforms (8 kDa, 12-19 kDa and 25-32 kDa) with long or short N-terminus and a great diversity in the exonic composition of C-terminus generated by alternative pre-mRNA splicing and differential usage of two translational start sites. RDM1 protein can be expressed differently in cytoplasm or nuclear/nucleolar in the absence of stress, proteotoxic stress and mild heat shock, suggesting functional differences among the RDM1 proteins. (PMID: 28939762, 17905820, 28939762, 30069034)

Notable Publications

Author	Pubmed ID	Journal	Application
Yajun Chen	31222930	J Cell Mol Med	WB
Zewei Tu	33634092	Front Cell Dev Biol	WB
Guojin Xie	30868057	FEBS Open Bio	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




PC-3 cells were subjected to SDS PAGE followed by western blot with 20156-1-AP (RDM1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 20156-1-AP (RDM1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 20156-1-AP (RDM1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed PC-3 cells using 20156-1-AP (RDM1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).