For Research Use Only

## ISCA1 Polyclonal antibody Catalog Number:20324-1-AP




| Basic Information      | Catalog Number:<br>20324-1-AP   | GenBank Accession Number:<br>BC002675                                   | Purification Method:<br>Antigen affinity purification |
|------------------------|---|---|---|
|                        | Size:<br>500 ug/ml  | GenelD (NCBI):<br>81689   | Recommended Dilutions:<br>WB 1:500-1:1000             |
|                        | Source:<br>Rabbit   | UNIPROT ID:<br>Q9BUE6   |   |
|                        | Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG14188   | Full Name:<br>iron-sulfur cluster assembly 1<br>homolog (S. cerevisiae) |   |
|                        |   | Calculated MW:<br>129 aa, 14 kDa  |   |
|                        |   | Observed MW:<br>15 kDa  |   |
| Applications           | Tested Applications:  | Positive C  | ontrols:  |
|                        | WB, ELISA<br>Species Specificity:<br>human, mouse   | WB : HepG2 cells, mouse cerebellum tissue, mouse heart tissue           |   |
| Background Information | ISCA1, also named as HBLD2 and hIscA, belongs to the hesB/iscA family. It is involved in the assembly of mitochondrial iron-sulfur proteins. ISCA1 probably involved in the binding of an intermediate of Fe/S cluster assembly. ISCA1 plays an important role in both mitochondrial and cytosolic iron-sulfur cluster biogenesis, and a notable component of the latter is the interaction between ISCA1 and IOP1. |   |   |
| Storage                | Storage:<br>Store at -20°C. Stable for one year<br>Storage Buffer:<br>PBS with 0.02% sodium azide and<br>Aliquoting is unnecessary for -20°   | i 50% glycerol pH 7.3.  |   |

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20324-1-AP (ISCA1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.