

For Research Use Only

PRKAR1A Polyclonal antibody

Catalog Number: 20358-1-AP

2 Publications



Basic Information

Catalog Number:

20358-1-AP

Size:

1000 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG14216

GenBank Accession Number:

BC036285

GeneID (NCBI):

5573

UNIPROT ID:

P10644

Full Name:

protein kinase, cAMP-dependent, regulatory, type I, alpha (tissue specific extinguisher 1)

Calculated MW:

381 aa, 43 kDa

Observed MW:

43-45 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IHC 1:250-1:1000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB : HeLa cells, mouse brain tissue, rat brain tissue

IHC : mouse testis tissue, human stomach cancer tissue, mouse brain tissue, rat salivary gland tissue

IF/ICC : HeLa cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

PKA is a heterotetramer inactive kinase composed of two regulatory and two catalytic subunits. The regulatory subunits are encoded by four genes (PRKAR1A, PRKAR2A, PRKAR1B, and PRKAR2B). PRKAR1A, a gene coding for the cAMP-dependent protein kinase (PKA) 1 α regulatory subunit, is located on human chromosome 17q22-24. PRKAR1A protein insufficiency and PKA dysregulation have been implicated in various types of disorders, including Albright hereditary osteodystrophy (AHO), pseudohypoparathyroidism (PHP), acrodysostosis (ACRDYS), and Carney complex. PRKAR1A protein expression level was significantly dysregulated in multiple primary carcinomas and distant metastases, such as cardiac myxomas, odontogenic myxomas, anaplastic thyroid carcinomas, breast cancer, pediatric pituitary adenomas, and Schwann cell tumors.

Notable Publications

Author	Pubmed ID	Journal	Application
Susanne Zellner	33545068	Mol Cell	WB
Yijia Ren	39341811	Nat Commun	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20358-1-AP (PRKAR1A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 20358-1-AP (PRKAR1A antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 20358-1-AP (PRKAR1A antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 20358-1-AP (PRKAR1A antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat salivary gland tissue slide using 20358-1-AP (PRKAR1A antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat salivary gland tissue slide using 20358-1-AP (PRKAR1A antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 20358-1-AP (PRKAR1A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 20358-1-AP (PRKAR1A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 20358-1-AP (PRKAR1A antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol)
fixed HeLa cells using PRKAR1A antibody (20358-1-
AP) at dilution of 1:400 and Multi-rAb CoraLite ®
Plus 488-Goat Anti-Rabbit Recombinant Secondary
Antibody (H+L) (RGAR002).