

For Research Use Only

STAC3 Polyclonal antibody

Catalog Number: 20392-1-AP

Featured Product

9 Publications



Basic Information

Catalog Number:

20392-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG14232

GenBank Accession Number:

BC008069

GeneID (NCBI):

246329

UNIPROT ID:

Q96MF2

Full Name:

SH3 and cysteine rich domain 3

Calculated MW:

364 aa, 42 kDa

Observed MW:

42-55 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:14000

IHC 1:20-1:200

IF-P 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF-P, ELISA

Cited Applications:

WB, IHC, IF, ColP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse skeletal muscle tissue, rat skeletal muscle tissue

IHC : human gliomas tissue, human heart tissue

IF-P : mouse skeletal muscle tissue,

Background Information

Notable Publications

| Author | Pubmed ID | Journal | Application |
|------------------|-----------|---------------------------|-------------|
| Yuki Ashida | 34590910 | J Appl Physiol (1985) | WB |
| Yuki Ashida | 35912995 | Am J Physiol Cell Physiol | WB |
| Yousra El Ghaleb | 35349630 | J Gen Physiol | WB |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20392-1-AP (STAC3 antibody) at dilution of 1:7000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human gliomas using 20392-1-AP (STAC3 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human gliomas using 20392-1-AP (STAC3 antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed mouse skeletal muscle tissue using STAC3 antibody (20392-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse skeletal muscle tissue using STAC3 antibody (20392-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).