For Research Use Only

CXCR7 Polyclonal antibody Catalog Number: 20423-1-AP Featured Product

Featured Product 22 Publications




Basic Information	Catalog Number: 20423-1-AP	GenBank Accession Num BC036661	ber: Purification Met Antigen affinity	
	Concentration:	GenelD (NCBI): 57007	Recommended I WB 1:500-1:1000	
	153 μg/ml Source: Rabbit	UNIPROT ID: P25106	IP 0.5-4.0 ug for protein lysate	, 1.0-3.0 mg of total
	Isotype:Full Name:IgGchemokine (C-X-C motif) receptor 7		IHC 1:50-1:500 receptor 7	
	Immunogen Catalog Number:Calculated MW:AG14247362 aa, 41 kDa			
		Observed MW: 45-50 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IP, ELISA Cited Applications:		WB : Raji cells, Jurkat cells, K-562 cells, mouse thymus tissue	
	WB, IHC, IF		IP : HUVEC cells,	
	Species Specificity: human, mouse, rat	IHC : mouse brain tissue,		
	Cited Species: human, mouse, rat			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	CXCR7 (C-X-C chemokine receptor family. CXCR7 can bind the chem CXCR4 for a restricted number of as well as with tumor growth and experimentally determined mole	okines CXCL11 and CXCL12 w HIV isolates. Expression of CX progression. This antibody re	ith high affinity, and it also ac CR7 has been associated with cognizes endogenous CXCR7,	ts as coreceptor with cardiac development
Notable Publications	Author	Pubmed ID Journal		Application
	Lulu Chen	32876991 Cell Biol	Int	WB,IF,IHC
	Liang Kuang	31662319 Ann Rhe	um Dis	WB
	Xiang Li	36405246 J Oncol		WB


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Raji cells were subjected to SDS PAGE followed by western blot with 20423-1-AP (CXCR7 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-CXCR7 (IP:20423-1-AP, 3ug; Detection:20423-1-AP 1:500) with HUVEC cells lysate 1400ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 20423-1-AP (CXCR7 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 20423-1-AP (CXCR7 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).