## For Research Use Only

## INSR/CD220 Polyclonal antibody

Catalog Number:20433-1-AP 31 Publications



**Basic Information** 

Catalog Number: 20433-1-AP Concentration:

500 ug/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG13922 1382 aa, 156 kDa

Observed MW:

Calculated MW:

190 kDa, 135 kDa, 95 kDa

GenBank Accession Number:

BC117172

GeneID (NCBI):

**UNIPROT ID:** 

P06213 Full Name:

INSR

**Purification Method:** Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

**Applications** 

**Tested Applications:** WB, IP, ELISA

Cited Applications: WB, IHC, IF, CoIP, Dot blot

Species Specificity:

human **Cited Species:** 

human, mouse, rat, pig, zebrafish, fish

Positive Controls:

WB: HEK-293 cells, MCF-7 cells, HT-1080 cells, HeLa

cells

IP: HeLa cells,

**Background Information** 

The biological effects of INS are mediated by a membrane-spanning cell surface receptor that has been shown to consist of disulfide-linked alpha-subunits (135 kDa) and beta-subunits (95 kDa) that are cleaved products of a common precursor. The binding of INS to the receptor extracellular domain results in intracellular activation of a tyrosine-specific kinase activity and the generation of signals that determine the cellular response (PMID: 2369896). This antibody raised against 980-1382aa of human INS receptor can recognize full-length INS receptor precursor and INS receptor beta subunit.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Takashi Hara	36130217	Biosci Biotechnol Biochem	WB
Yu Bai	34603025	Front Pharmacol	WB,CoIP
Si-Lan Han	33245450	Fish Physiol Biochem	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data





IP result of anti-INSR (IP:20433-1-AP, 4ug; Detection:20433-1-AP 1:500) with HeLa cells lysate 1600ug.

HEK-293 cells were subjected to SDS PAGE followed by western blot with 20433-1-AP (INSR antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.