For Research Use Only

SLC37A2 Polyclonal antibody

Catalog Number:20469-1-AP 1 Publications



Basic Information

Catalog Number: 20469-1-AP

Size: 500 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG14306

GenBank Accession Number:

BC051314 GeneID (NCBI): 219855 UNIPROT ID: Q8TED4

solute carrier family 37 (glycerol-3phosphate transporter), member 2

Calculated MW: 505 aa, 55 kDa Observed MW: 50-75 kDa

Full Name:

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC

Species Specificity: human, mouse, rat Cited Species: human Positive Controls:

WB: mouse thymus tissue, HeLa cells, RAW 264.7 cells, mouse spleen tissue, rat thymus tissue

Purification Method:

WB 1:300-1:600

Antigen affinity purification

Recommended Dilutions:

Background Information

The solute carrier family 37 (SLC37) consists of four sugar-phosphate exchangers, A1, A2, A3, and A4, which are anchored in the endoplasmic reticulum (ER) membrane (PMID:23506893). Solute Carrier Family 37 Member 2 (SLC37A2), also knows as SPX2, was firstly identified in a work, conducted on mice and aimed to detect cAMP inducible genes playing a role in promoting cholesterol efflux from the macrophage cell line RAW264 via apoE and apoA1 (PMID:11004510). Interestingly, of the four SLC37A members, SLC37A2 displays the highest level of transcript abundance in neutrophils and macrophages, indicating that SLC37A2 may play an essential role in regulating innate immune function (PMID:32428862).

Notable Publications

Author	Pubmed ID	Journal	Application
Xinwei Xu	39259386	Cell Biol Toxicol	WB,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



mouse thymus tissue were subjected to SDS PAGE followed by western blot with 20469-1-AP (SLC37A2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 20469-1-AP (SLC37A2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



mouse spleen tissue were subjected to SDS PAGE followed by western blot with 20469-1-AP (SLC37A2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.