

PI3 Kinase p110 Alpha Polyclonal antibody

Catalog Number: 20583-1-AP

12 Publications

Basic Information

Catalog Number:

20583-1-AP

Size:

193 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_006218

GeneID (NCBI):

5290

UNIPROT ID:

P42336

Full Name:

phosphoinositide-3-kinase, catalytic,
alpha polypeptide

Calculated MW:

124 kDa

Observed MW:

120-130 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:300-1:1000

IHC 1:50-1:500

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HepG2 cells, COLO 320 cells, HeLa cells, A431 cells, mouse liver tissue

IHC : human cervical cancer tissue,

Background Information

PIK3CA belongs to the PI3/PI4-kinase family. It phosphorylates PtdIns, PtdIns4P and PtdIns(4,5)P2 with a preference for PtdIns(4,5)P2. Defects in PIK3CA are associated with colorectal cancer (CRC). Defects in PIK3CA are associated with breast cancer. Defects in PIK3CA are associated with ovarian cancer. Defects in PIK3CA may underlie hepatocellular carcinoma (HCC). Defects in PIK3CA are a cause of keratosis seborrheic (KERSEB). The antibody is specific to PIK3CA.

Notable Publications

Author	Pubmed ID	Journal	Application
Wu Jun J	23582786	Biomed Pharmacother	WB
H Zhang	28581513	Oncogene	WB,IHC
Hua Wang	27326255	J Cancer	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 20583-1-AP (PI3K p110(alpha) antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human cervical cancer using 20583-1-AP (PI3K p110(alpha) antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded human cervical cancer using 20583-1-AP (PI3K p110(alpha) antibody) at dilution of 1:50 (under 10x lens).



HeLa cells were subjected to SDS PAGE followed by western blot with 20583-1-AP (PI3K p110(alpha) antibody at dilution of 1:300 incubated at room temperature for 1.5 hours.