

For Research Use Only

CXCR2 Polyclonal antibody, PBS Only

Catalog Number: 20634-1-PBS

Featured Product



Basic Information

Catalog Number:

20634-1-PBS

Concentration:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG14670

GenBank Accession Number:

BC037961

GeneID (NCBI):

3579

UNIPROT ID:

P25025

Full Name:

interleukin 8 receptor, beta

Calculated MW:

360 aa, 41 kDa

Observed MW:

45-50 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IHC, IF-P, IP, ELISA

Species Specificity:

human

Background Information

IL8RB, also named as CXCR2, GRO/MGSA receptor, IL-8 receptor type 2, CDw128b, and CD182, belongs to the G-protein coupled receptor 1 family. It is a receptor for interleukin-8 which is a powerful neutrophil chemotactic factor. The binding of IL-8 to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. It binds to IL-8 with high affinity. IL8RB also binds with high affinity to CXCL3, GRO/MGSA, and NAP-2.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




L02 cells were subjected to SDS PAGE followed by western blot with 20634-1-AP (CXCR2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 20634-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 20634-1-AP (CXCR2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 20634-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 20634-1-AP (CXCR2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 20634-1-PBS in a different storage buffer formulation.



IP result of anti-CXCR2 (IP:20634-1-AP, 4ug; Detection:20634-1-AP 1:1000) with HepG2 cells lysate 3200ug. This data was developed using the same antibody clone with 20634-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse liver tissue using CXCR2 antibody (20634-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 20634-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CXCR2 antibody (20634-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 20634-1-PBS in a different storage buffer formulation.