For Research Use Only

BRCA1 Polyclonal antibody

Catalog Number:20649-1-AP

16 Publications



Basic Information	Catalog Number:GenBank Accession Number:Purification Method:20649-1-APNM_007294Antigen affinity purification				
	Size: GeneID (NCBI): 390 μg/ml 672				
	Source: UNIPROTID: Rabbit P38398				
	Isotype: Full Name: IgG breast cancer 1, early onset				
	Calculated MW: 208 kDa Observed MW: 230-260 kDa				
Applications	Tested Applications: ELISA				
	Cited Applications: WB, IF, IHC				
	Species Specificity: human, mouse, rat				
	Cited Species: human, rat, mouse				
Background Information	RCA1, also named as RNF53, plays a central role in DNA repair by facilitating cellular response to DNA repair. It is required for appropriate cell cycle arrests after ionizing irradiation in both the S-phase and the G2 phase of the cell cycle. The BRCA1-BARD1 heterodimer coordinates a diverse range of cellular pathways such as DNA damage repai ubiquitination and transcriptional regulation to maintain genomic stability. BRCA1 acts by mediating ubiquitin E3 ligase activity that is required for its tumor suppressor function. It is involved in transcriptional regulation of P21 i response to DNA damage. BRCA1 inhibits lipid synthesis by binding to inactive phosphorylated ACACA and preventing its dephosphorylation. The antibody is specific to BRCA1 appears to produce multiple splice variants. BRCA1 is a nuclear protein with a molecular mass of 220 kDa. The present study describes the isolation a expression of two cDNAs of BRCA1, including a splice variant designated BRCA1D672-4095. BRCA1D672-4095 is generated by exclusion of exon 11 by in-frame splicing and produces a 97 kDa protein. In contrast to BRCA1, BRCA10672-4095 localizes to the cytoplasm.				
	transcriptional regulator. B preventing its dephosphory variants. BRCA1 is a nuclea expression of two cDNAs of generated by exclusion of	RCA1 inhibits lipid synth lation. The antibody is s r protein with a molecul BRCA1, including a split exon 11 by in-frame split	pecific to BRCA1. BR ar mass of 220 kDa. ce variant designate	CA1 appears to produce of the present study described BRCA1D672-4095. BRC	CACA and multiple splice pes the isolation a A1D672-4095 is
Notable Publications	transcriptional regulator. B preventing its dephosphory variants. BRCA1 is a nuclea expression of two cDNAs of generated by exclusion of BRCA1D672-4095 localizes	RCA1 inhibits lipid synth lation. The antibody is s r protein with a molecul BRCA1, including a spli exon 11 by in-frame split to the cytoplasm.	pecific to BRCA1. BR ar mass of 220 kDa. ce variant designate cing and produces a	CA1 appears to produce of the present study described BRCA1D672-4095. BRC	CACA and multiple splice bes the isolation a A1D672-4095 is st to BRCA1,
Notable Publications	transcriptional regulator. B preventing its dephosphory variants. BRCA1 is a nuclea expression of two cDNAs of generated by exclusion of BRCA1D672-4095 localizes	RCA1 inhibits lipid synth (lation. The antibody is s r protein with a molecul (BRCA1, including a splitexon 11 by in-frame splite to the cytoplasm.	pecific to BRCA1. BR ar mass of 220 kDa. ce variant designate cing and produces a Journal	CA1 appears to produce of the present study described BRCA1D672-4095. BRC	CACA and multiple splice bes the isolation a A1D672-4095 is st to BRCA1, Application
Notable Publications	transcriptional regulator. B preventing its dephosphory variants. BRCA1 is a nuclea expression of two cDNAs of generated by exclusion of G BRCA1D672-4095 localizes Author Yang Yu	RCA1 inhibits lipid synth lation. The antibody is s r protein with a molecul BRCA1, including a split exon 11 by in-frame split to the cytoplasm. Pubmed ID 31636387	pecific to BRCA1. BR ar mass of 220 kDa. ce variant designate cing and produces a Journal Oncogene	CA1 appears to produce of the present study described BRCA1D672-4095. BRC	CACA and multiple splice bes the isolation a A1D672-4095 is st to BRCA1,
Notable Publications	transcriptional regulator. B preventing its dephosphory variants. BRCA1 is a nuclea expression of two cDNAs of generated by exclusion of BRCA1D672-4095 localizes	RCA1 inhibits lipid synth (lation. The antibody is s r protein with a molecul (BRCA1, including a splitexon 11 by in-frame splite to the cytoplasm.	pecific to BRCA1. BR ar mass of 220 kDa. ce variant designate cing and produces a Journal	CA1 appears to produce of the present study described BRCA1D672-4095. BRC	CACA and multiple splice bes the isolation a A1D672-4095 is st to BRCA1, Application IF

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer. Selected Validation Data