

For Research Use Only

MARCKS Polyclonal antibody

Catalog Number: 20661-1-AP

5 Publications



Basic Information

Catalog Number:

20661-1-AP

Size:

300 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_002356

GeneID (NCBI):

4082

UNIPROT ID:

P29966

Full Name:

myristoylated alanine-rich protein kinase C substrate

Calculated MW:

32 kDa

Observed MW:

80 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF-P 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF-P, IP, ELISA

Cited Applications:

WB, IF, IP

Species Specificity:

human, mouse, rat

Cited Species:

human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HEK-293 cells, IMR-32 cells, mouse brain tissue, rat brain tissue, SH-SY5Y cells, SK-N-SH cells

IP : A431 cells,

IHC : human normal colon,

IF-P : mouse brain tissue,

Background Information

MARCKS, also named as MACS, PRKCSL and 80K-L protein, belongs to the MARCKS family. MARCKS is the most prominent cellular substrate for protein kinase C. It binds calmodulin, actin, and synapsin. MARCKS is a filamentous (F) actin cross-linking protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Nicholas J Eustace	30942445	Int J Oncol	IP
Falguni Das	35835217	J Biol Chem	WB
Tong Lei	32800794	J Proteomics	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20661-1-AP (MARCKS antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human normal colon slide using 20661-1-AP (MARCKS antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using MARCKS antibody (20661-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).



IP result of anti-MARCKS (IP:20661-1-AP, 4ug; Detection:20661-1-AP 1:300) with A431 cells lysate 1200ug.



Various lysates were subjected to SDS PAGE followed by western blot with 20661-1-AP (MARCKS antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.