

For Research Use Only

NPT1 Polyclonal antibody

Catalog Number: 20751-1-AP

3 Publications



Basic Information

Catalog Number:

20751-1-AP

Concentration:

400 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG12819

GenBank Accession Number:

BC101745

GeneID (NCBI):

6568

UNIPROT ID:

Q14916

Full Name:

solute carrier family 17 (sodium phosphate), member 1

Calculated MW:

467 aa, 51 kDa

Observed MW:

50 kDa, 60-64 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB : mouse kidney tissue, human placenta tissue

Background Information

NPT1 (also known as SLC17A1) is a member of sodium-dependent inorganic phosphate transporters (NPT), which regulates entrance into the cellular membrane. NPT1 is mainly expressed in the kidney transporting small organic anions such as PAH (para-aminohippurate), but it is also found in the liver and brain. NPT1 localizes to the apical membrane of renal proximal tubular cells. NPT1 exists two isoforms due to the alternative splicing. Western blot analysis using this antibody detected a major band around 45-55 kDa in kidney tissue.

Notable Publications

Author	Pubmed ID	Journal	Application
Ruixia Bao	34785103	Phytomedicine	WB
Jing Xiao	29497172	Exp Mol Med	WB
Ruixia Bao	31920654	Front Pharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 20751-1-AP (NPT1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.