

For Research Use Only

# MAP7D1 Polyclonal antibody, PBS Only

Catalog Number:20788-1-PBS



## Basic Information

<b>Catalog Number:</b> 20788-1-PBS	<b>GenBank Accession Number:</b> BC003083	<b>Purification Method:</b> Antigen affinity purification
<b>Concentration:</b> 1 mg/ml	<b>GeneID (NCBI):</b> 55700	
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q3KQU3	
<b>Isotype:</b> IgG	<b>Full Name:</b> MAP7 domain containing 1	
<b>Immunogen Catalog Number:</b> AG14385	<b>Calculated MW:</b> 841 aa, 93 kDa	
	<b>Observed MW:</b> 120-130 kDa	

## Applications

**Tested Applications:**  
WB, IHC, IF/ICC, IP, Indirect ELISA

**Species Specificity:**  
human

## Background Information

MAP7D1 also known as RPRC1, PARCC1, belongs to the MAP7 family. The MAP7 (Microtubule Associated Protein 7) protein family, consisting of four members, MAP7, MAP7D1, and MAP7D2, MAP7D3, is the microtubule-associated protein involved in various cellular processes regulating microtubule dynamics, organization, and stability (PMID: 28980356). MAP7D1 exhibits the highest conservation with MAP7 and was recently identified as a phosphorylation substrate of DCLK1 in cortical neurons. MAP7D1 is required to maintain MT acetylation, which is enriched in stable MTs (PMID: 35470240). Consistent with the literature, the apparent molecular mass of MAP7D1 detected by Western blot was 120-130 kDa (PMID: 35470240, 37550720).

## Storage

**Storage:**  
Store at -80°C.  
**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

**Storage Buffer:**  
PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20788-1-AP (MAP7D1 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 20788-1-PBS in a different storage buffer formulation.




IP result of anti-MAP7D1 (IP:20788-1-AP, 4ug; Detection:20788-1-AP 1:30000) with HEK-293 cells lysate 1470 ug. This data was developed using the same antibody clone with 20788-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 20788-1-AP (MAP7D1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 20788-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 20788-1-AP (MAP7D1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 20788-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed U-251 cells using MAP7D1 antibody (20788-1-AP) at dilution of 1:200 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-Phalloidin (red). This data was developed using the same antibody clone with 20788-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MAP7D1 antibody (20788-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 20788-1-PBS in a different storage buffer formulation.